



Simon Baron-Cohen, Bonnie Auyeung, Emma Ashwin, Rebecca Knickmeyer

The Extreme Male Brain Theory of Autism:

3 The Role of Fetal Androgens

.

10

12

13

14

15

16

17

18

19

20

Points of Interest

- There are sex differences in neuroanatomy, neural function, cognition, and behavior in the general population.
- People with ASD show an extreme of the typical male profile in terms of empathy and systemizing.
- Whether such hyper-masculinization is evident at the level of neuroanatomy and neural function in ASD remains to be tested.
- Fetal testosterone (fT) is known from animal research to play an organizing role in brain development.
- fT associated with individual differences in eye contact, vocabulary development, empathy, systemizing, attention to detail, and autistic traits in typically developing children.
- It remains to be tested whether fT is elevated in children who go on to develop an ASD.

Autism and Asperger Syndrome (AS) are autism spectrum disorders (ASDs). The diagnosis rests on the presence of difficulties in reciprocal social interaction and communication, along with strongly repetitive behavior and unusually narrow interests (A.P.A., 1994). The prevalence of ASD is currently estimated to be 1% (Scott et al., 2002). Autism spectrum disorder is biased toward males (Fombonne, 2005) with a 4:1 (male:female) ratio for classic autism (Chakrabarti & Fombonne, 2001) and 8:1 for AS (Scott et al., 2002). Classic autism and AS differ in terms of the presence of additional learning difficulties and language delay (in classic autism 31 only). Autism spectrum disorders are neurobiological and genetic (Stodgell et al., 2001), but the specific factors responsible for the higher male incidence in ASDs remain unclear. One possible factor is that ASDs may be an extreme of the male brain (Baron-Cohen, 2002).

iii

The Extreme Male Brain: Psychology

The extreme male brain (EMB) theory of autism is an extension 39 of the empathizing-systemizing (E-S) theory of typical sex 40 differences (Baron-Cohen, 2003). The latter theory proposes 41 that females, on average, have a stronger drive to empathize 42 (to identify another person's thoughts and feelings and to 43 respond to these with an appropriate emotion), whereas males, 44 on average, have a stronger drive to systemize (to analyze or 45 construct rule-based systems) (Baron-Cohen, 2003). Evidence 46 relevant to this is as follows: Individuals with ASD score higher 47 on the systemizing quotient (SQ), an instrument on which 48 typical males score higher than typical females (Baron-Cohen 49 et al., 2003; Auyeung et al., 2006; Wheelwright et al., 2006). 50 Individuals with ASDs have intact or superior functioning on 51 tests of intuitive physics (Baron-Cohen et al., 2001; Lawson 52 et al., 2004), a domain that shows a sex difference in favor of 53 males (Lawson et al., 2004). Individuals with ASDs are faster 54 and more accurate than controls on the Embedded Figures 55 Task (EFT), a task on which typical males perform better than 56 typical females (Shah & Frith, 1983; Jolliffe & Baron-Cohen, 57 1997). The EFT requires good attention to detail, a prerequisite for systemizing. All of these are relevant to systemizing.

On the EQ (Baron-Cohen & Wheelwright, 2004), individuals 60 with ASDs score lower than control groups, whereas typical 61 females score higher than typical males (Baron-Cohen & 62 Wheelwright, 2004). Individuals with ASDs score lower than 63 typical males on the "Reading the Mind in the Eyes" task 64 (Baron-Cohen et al., 1997), the Social Stories Questionnaire 65 (Lawson et al., 2004), on tests of recognizing complex emotions 66 from videos of facial expressions or audios of vocalizations 67 (Golan et al., 2006), and on the Friendship and Relationship 68 Questionnaire (which tests the importance of emotional 69 intimacy and sharing in relationships) (Baron-Cohen & 70

Insert
Empathizing

Empathizing
Quotient; put
EQ in
parentheses





989



990 Autism Spectrum Disorders

1 Wheelwright, 2003). All of these are tests of empathy. On the Childhood Autism Spectrum Test (CAST) (Scott et al., 2002; Scott et al., 2002), boys score higher than girls (Williams et al., 2008), and children with ASDs score higher than controls (Williams et al., 2005). On the autism spectrum quotient (AQ) (Baron-Cohen et al., 2001a), individuals with ASDs score higher than those without a diagnosis (Baron-Cohen et al., 2001). Among controls, males score higher than females (Baron-Cohen et al., 2001; Baron-Cohen et al., 2006; Auyeung et al., 2008), a finding that has been reported cross-culturally (Wakabayashi et al., 2004; Wakabayashi et al., 2006; 11 Wakabayashi et al., 2007; Hoekstra et al., 2008). Similar results 12 have been found using the Social Responsiveness Scale (SRS), finding that individuals with an ASD diagnosis score higher than typical males, who in turn score higher than typical females (Constantino & Todd, 2005). All of these are measures of autistic traits.

18

19

The Extreme Male Brain: Biology

Characteristics of neurodevelopment in autism may also 20 represent an exaggeration of typical sex differences in brain 21 development (Baron-Cohen et al., 2005). For example, there 22 is larger overall brain volume in males during childhood 23 (Giedd et al., 1996), and in ASDs, this may be even more 24 extreme (Courchesne et al., 2004). There is greater growth of the amygdala in males during childhood (Merke et al., 2005), and in ASDs, this may be even more extreme (Sparks et al., 2002). The posterior section of the corpus callosum is thicker in females than males (Jancke et al., 1997) and in ASDs is even 29 thinner than is typical (Herbert et al., 2004). These are all volumetric measures of the brain. Using fMRI, typical females show increased activity in the extrastriate cortex during the 32 EFT and increased activity bilaterally in the inferior frontal 33 cortex during the "Reading the Mind in the Eyes" Test 34 compared to typical males (Baron-Cohen et al., 2006). People 35 with ASDs show even less activity in each of these regions 36 during these tasks (Baron-Cohen et al., 1999; Ring et al., 1999). Parents of children with ASDs also show hyper-masculinization of brain activity (Baron-Cohen et al., 2006), suggesting that 39 this may be part of the broader autism phenotype. These are all functional neuroimaging measures of the brain. It remains important to identify the biological mechanisms that cause such sexual dimorphism. One possible biological mechanism is the effect of fetal testosterone (fT) (Geschwind & Galaburda, 1985), reviewed in the next section.

46

The Role of Fetal Testosterone in Brain Development

Fetal gonadal hormones (including the androgens [e.g., testosterone, dihydrotestosterone], estrogens [e.g., estradiol,




Figure 55–1. The Empathizing-Systemizing Model of Typical Sex Differences. The Main Brain Types are Illustrated on Axes of Empathising (E) and Systemising (S) Dimensions (Numbers Represent Standard Deviations from the Mean). Balanced Brain (Type B, Grey Region); Female Brain (Type E, Light Blue Region), Male Brain (Type S, Red Region); the Extreme Types E (Dark Blue) and S (Dark Red) Lie at the Outer Borders. According to the "Extreme Male Brain" theory of Autism, People with ASD will Generally Fall in the Dark Red Region. Modified with Permission from Baron-Cohen S. (2002) The Extreme Male Brain Theory of Autism. *Trends in Cognitive Science* 6, 248–254.

Extreme S

estrone, estriol], and progestins [e.g., progesterone]) lead to the differentiation of the male and female phenotype (Fuchs & Klopper, 1983; Tulchinsky & Little, 1994; Kimura, 1999; Hines, 2004). If androgens and the androgen receptors are present, the male genital phenotype will develop. If not (as seen in Complete Androgen Insensitivity Syndrome), the female genital phenotype will develop (Jost, 1961; Jost, 1970, 1972; George et al., 1992). Fetal gonadal hormones are essential for sexual differentiation of both the body and the brain (Goy, 1980; Fitch & Denenberg, 1998). In what follows, we summarize some key points in this process. Three surges in testosterone levels are known to occur. The first surge is between weeks 8 and 24 of gestation (Collaer & Hines, 1995; Baron-Cohen et al., 2004; Hines, 2004). Most of the prenatal androgen effects occur between 7 and 12 weeks of gestation (Rommerts, 2001). Then, after birth, a second peak in circulating testosterone occurs in human male infants. Usually the levels remain in the pubertal range for a few months and then drop to the barely detectable levels observed in childhood by age 4 to 6 months (Smail et al., 1981). Finally, the third surge is associated with puberty. Individuals vary both in the levels of hormones to which they are exposed and in their sensitivity



51

52

53

54

55

57

58

59

60

61

63

64

65

66

67

69

70

71



1 to those hormones. Variations in androgen responsivity (caused by mutations in the human androgen receptor gene) can result in complete insensitivity to androgens (and thus female differentiation) or to infertility and minor undervirilization (Casella et al., 2001). The timing of hormonal effects influences whether effects are organizational or activational (Goy, 1980). Organizational effects produce permanent changes in the brain (Phoenix et al., 1959) and are most likely to occur during early development when most neural structures are becoming established. The discovery of such organizational effects of fT came from animal research (mostly rodents), and to date, these have only looked at neuroanatomy or limited aspects of behavior (spatial ability and mating). Activational effects occur later and are associated with concurrent changes in circulating hormone levels (Kimura & Hampson, 1994; Cooke et al., 1999).

Fetal Androgens Affect Brain and Behavior: Evidence from Rare Medical Conditions

Although animal research provides strong evidence for the 20 effect of hormones in development, the direct manipulation of hormones in human fetuses is unethical. Instead, researchers have studied people, in whom for medical reasons the sex hormones are higher or lower than expected for a person's sex (Money & Ehrhardt, 1972). Two examples are Congenital Adrenal Hyperplasia (CAH) and Complete Androgen 26 Insensitivity Syndrome (CAIS). Congenital Adrenal Hyperplasia 27 is a condition in which an enzymatic defect (usually caused by 28 mutations in the gene coding for 21-hydroxylase) results 29 in high levels of adrenal androgens, beginning very early in gestation. It has an incidence of 1 in 10,000 to 1 in 15,000 live 31 births (Grumbach et al., 2003). Females with CAH differ from 32 unaffected females (their siblings or age- and sex-matched 33 controls) in spatial orienting, visualization, and targeting 34 (Resnick et al., 1986; Hampson et al., 1998; Hines et al., 2003b). 35 Females with CAH are also more likely than controls to be left-handed (Nass et al., 1987), although the difference is small and inconsistent. Females with CAH are more interested in male-typical activities and less interested in female-typical 39 activities (Ehrhardt & Baker, 1974; Berenbaum & Hines, 1992; Berenbaum & Snyder, 1995; Berenbaum, 1999; Hines et al., 2004). Girls with CAH score lower than sex-matched controls on measures assessing empathy, intimacy, and the need for close social relationships (Mallin & Walker, 1972; Grimshaw et al., 1995). Individuals with CAH showed higher levels of language and learning difficulties than unaffected family members (Resnick et al., 1986). All these changes can 47 be interpreted as masculinization. 48

Complete Androgen Insensitivity Syndrome is a second example of a rare medical condition affecting prenatal endocrine responses. It occurs when there is a complete deficiency of working androgen receptors. It is an X-linked recessive disorder and hence occurs more often in genetic males.

References are incorrect on lines 44 and 45: should be Mathews et al (2009), Helleday et al (1993), Resnick (1982) and Kuhnle & Bullinger (1997)

Prevalence is between 1 in 20,000 and 1 in 60,000 live male 54 births. At birth, genetic male infants with CAIS are phenotypically female, despite an XY complement, and are usually 56 raised as girls. At puberty, breasts develop under the influence of estrogen derived from testicular androgens. Diagnosis usually takes place when menarche fails to occur (Nordenstrom 59 et al., 2002; Grumbach et al., 2003). In studies, gender identity 60 has not differed between genetic males with CAIS and control 61 women (Quadagno et al., 1977; Hines et al., 2003a). Individuals 62 with CAIS perform in a female-typical fashion on tests of 63 visuo-spatial ability (Money et al., 1984).

These findings suggest that two X chromosomes or functioning ovaries are not required for feminine-typical psychological development in humans and highlight the role of androgen receptors in influencing masculine-typical psychological development. However, existing studies have not examined all sexually dimorphic aspects of neurobiology and cognition in these populations. Therefore, one cannot rule out an effect of X-chromosome genes or ovarian hormones on some aspects of sex-typical neurodevelopment.

Fetal Androgens Affect Brain and Behavior: Evidence from Amniotic Fluid Testosterone

Another approach to testing if fetal androgens affect brain 77 and behavior is to measure fT in amniotic fluid obtained 78 during routine diagnostic amniocentesis. An advantage of this 79 approach is its timing. It is typically performed during the 80 second trimester of pregnancy (usually 14–20 weeks of gesta- 81 tion) that coincides with the serum testosterone peak period 82 in male fetuses. Several studies have documented a large sex 83 difference in amniotic androgen levels (Judd et al., 1976; 84 Dawood, 1977; Robinson et al., 1977; Nagami et al., 1979; 85 Finegan et al., 1989). The origin of androgens in amniotic 86 fluid is the fetus itself. Hormones enter the amniotic fluid via 87 diffusion through the fetal skin in early pregnancy and via fetal 88 urine in later pregnancy (Judd et al., 1976; Schindler, 1982). 89 Testosterone obtained in amniotic fluid is thought to be a 90 good reflection of the levels in the fetus (van de Beek et al., 91 2004) and represents an alternative to direct assay of fetal 92 serum that would be unnecessarily invasive. In the Cambridge 93 Fetal Androgen Project, children whose mothers had amnio- 94 centesis during pregnancy (but who were otherwise typically 95 developing children) were followed-up after birth at ages 12, 96 18, 24, 48, and 96 months (Baron-Cohen et al., 2004). Evidence 97 that amniotic testosterone affects cognitive development 98 includes the following. Fetal testosterone is inversely associ- 99 ated with eye contact in males at age 12 months (Lutchmaya 100 et al., 2002). Fetal testosterone is inversely associated with 101 size of vocabulary development at ages 18 and 24 months 102 (Lutchmaya et al., 2002). Fetal testosterone is inversely 103 associated with quality of social relationships, and posi- 104 tively associated with narrow interests, at age 48 months 105 (Knickmeyer et al., 2005). Fetal testosterone is inversely 106





74

75



992 **Autism Spectrum Disorders**

AU: Please update details.

Should read hypermasculinized associated with empathy at ages 48 and 96 months (Chapman et al., 2006; Knickmeyer et al., 2006). Fetal testosterone is positively associated with "systemizing" at age 96 months (Auyeung et al., 2006). Fetal testosterone is positively associated with performance on the EFT, as a measure of attention to detail, at age 96 months (Auyeung et al., submitted). The effect sizes in these studies range from 0.2 to 0.4. Because all of these domains of behavior (eye contact, language development, quality of social relationships, narrow interests, empathy, systemizing, and embedded figures/attention to detail) show sexual dimorphism and may be hyper-mASDulinized in ASDs, it raises the question as to whether fetal testosterone plays a role in the development of autism. In the final section, we review evidence that androgens—especially fetal testosterone—may play a role in autism or autistic traits.



17

20

21

22

26

27

28

29

31

32

33

34

35

38

39

40

41

42

44

45

46

47

The Role of Sex Steroid Hormones in Autism Spectrum Disorders or Autistic Traits

Evidence that supports a role for hormones in the development of ASDs is preliminary and includes the following: Androgen-related medical conditions such as polycystic ovary syndrome (PCOS), ovarian growths, and hirsutism occur with elevated rates in women with AS and in mothers of children with autism (Ingudomnukul et al., 2007). Girls with CAH have a higher AQ score than their unaffected sisters (Knickmeyer et al., 2006c). Children with AS and children with classic autism have lower 2D:4D ratios than typical developing children (Manning et al., 2001; Milne et al., 2006). The ratio of the second digit to the fourth digit (2D:4D ratio) is lower in men than in women. The 2D:4D ratio is fixed by week 14 of fetal life (Garn et al., 1975) and is influenced by testosterone (Manning et al., 2002). This suggests children with ASD have been exposed to higher amounts of androgens. A subset of male adolescents with autism show hyper-androgeny, or elevated levels of androgens, and precocious puberty (Tordjman et al., 1997). Delayed menarche has also been found in females with AS (Knickmeyer et al., 2006; Ingudomnukul et al., 2007). Puberty timing reflects hormonal programming of the hypothalamic-pituitary-gonadal axis during gestation (Grumbach & Shaw, 1998).

Left-handedness and ambidexterity are more common in typical males (Peters, 1991) and individuals with autism (Gillberg, 1983). Body asymmetries are related to prenatal sex hormones and breast or testis size on the left versus right sides of the body are related to cognition (Kimura, 1999). Fetal testosterone is implicated in left-handedness and asymmetric lateralization (Fein et al., 1985; Satz et al., 1985; Soper et al., 1986; McManus et al., 1992). The typical male brain is heavier than the female brain, a difference that partly results from fT exposure (Hines, 2004). Individuals with autism have even heavier brains than typical males (Hardan et al., 2001). Amniotic fT levels are positively associated with higher scores (indicating greater number of autistic traits) on the CAST and 54 on the child autism spectrum quotient (AQ-C) (Auyeung et al., 2008). These findings are consistent with the fetal androgen theory of autism, although the ultimate test of this theory 57 will require testing between fetal testosterone and clinically 58 diagnosed ASDs. The latter will require much larger samples 59 than have previously been tested.

It is important in a review chapter such as this to summarize 61 criticisms of the fetal androgen theory of ASDs. First, if autism 62 is an extreme of the typical male behavioral profile, why do people with ASD not show high levels of aggression or a strong interest in competitive sports? We suspect this criticism reflects a misunderstanding of the different roles that 66 fetal as opposed to current (circulating) testosterone play. 67 The latter may well affect aggression and competitiveness, but 68 fetal androgens may selectively affect very different aspects of 69 cognition such as attention to detail and empathy. This 70 remains to be tested. Second, given that testosterone interacts 71 with multiple systems in the body, why single it out for a 72 special role? We acknowledge that if fT plays a role in ASDs, 73 it is likely to do this in complex ways, as testosterone 74 modulates neurotransmitters (such as GABA) as well as peptide hormones (such as oxytocin), to name just two examples. Understanding such relationships will require testing of multiple systems within the same experiment. Third, given 78 that amniotic testosterone is currently only studied in humans 79 via amniocentesis and only 6% of pregnant women undergo 80 amniocentesis, does this not lead to potentially biased samples? We acknowledge this bias but would note that currently 82 this is the only ethical way to study fT because amniocentesis 83 itself carries a risk of inducing miscarriage (in 2% of cases) and so cannot be justified purely for research on a randomly selected, and therefore representative, sample of pregnant women. The main risk of bias comes from higher maternal age because this is one reason why women are referred for an amniocentesis (being older than 35 years old). For this reason, maternal age is entered as a variable in the regression analysis. 90 All fT effects that have been found remain significant after 91 removing any variance because of maternal age.

A further criticism is that if sex differences in the population 93 are only found when one compares equal numbers of males 94 and females, and if studies of ASDs tend to be biased toward males, is the extreme male brain theory really just a reflection that ASDs affect more males than females? This is a valid concern, but it interesting that where it is possible to compare equal numbers of males and females with ASDs, typical sex differences are absent, which argues against any risk of circu- 100 larity. Also, when fT effects are found on behavior, these are seen within sex, not just when the sexes are combined. This suggests these are hormone effects rather than a redescription 103 of sex differences. Finally, Skuse and others have pointed out 104 that sex-linked neural and behavioral phenotypes could emerge not just because of hormone effects but genetic effects. 106 We agree that fT is only one of many mechanisms that are 107 likely candidates for giving rise to such sex differences, and indeed our recent candidate gene study identified 10 genes 109

insert is

92

97





The Extreme Male Brain Theory of Autism: The Role of Fetal Androgens

- 1 involved in the sex steroid hormone pathway that were
- 2 nominally associated with either autistic traits, empathy, or an
- 3 ASD (Chakrabarti et al., 2009).

Conclusions

- 6 The higher incidence of autism in male individuals might provide important clues to the etiology of the condition, which
- has been described as an "extreme of the male brain" (Baron-
- Cohen, 2002). The studies reviewed here suggest that prenatal
- testosterone could be involved in the sex differences in key
- areas of behavior in the general population (social development,
- language development, empathy, systemizing, and attention
- to detail) and to the male vulnerability to autism. These studies
- suggest that variations in fetal testosterone are related to
- individual differences in cognition and behavior in typically
- developing children, but caution needs to be taken when
- extrapolating these results to individuals with autism. Our
- ongoing collaboration with the biobank in Denmark that has
- thousands of amniotic samples will enable a test of the fetal
- testosterone theory in clinically diagnosed cases of ASD.

AU: Update

"submitted"

details for

the

Challenges and Future Directions



25

26

27

28

29

30

31

32

33

34

- · Testing the fetal testosterone theory in relation to diagnosed ASD will require tens of thousands of amniotic samples because only 1 in 100 of these are expected to go on to develop an ASD.
- It is an assumption that amniotic testosterone reflects testosterone levels in the brain, but this remains untested.
- · Ethically, if it were established that in ASD there are elevated fetal testosterone levels, this does not mean a treatment implication is to block fT, as fT is likely to be involved in many systems, not just the development of autistic traits.

SUGGESTED READINGS

- Baron-Cohen S. (2002) The extreme male brain theory of autism. 36 37 Trends in Cognitive Science 6, 248-254.
- Baron-Cohen S, Lutchmaya S, & Knickmeyer R. (2004) Prenatal 38 testosterone in mind: Amniotic fluid studies. Cambridge, MA: 39 MIT/Bradford Books. 40
- Baron-Cohen S, Knickmeyer R, & Belmonte MK. (2005) Sex differ-41 ences in the brain: Implications for explaining autism. Science 42 310, 819-823. 43

ACKNOWLEDGMENTS

- SBC was supported by grants from the Nancy Lurie Marks Family
- Foundation and the MRC UK during the period of this work. We are

grateful to Svetlana Lutchmaya, Bhismadev Chakrabarti, and Mike 47 Lombardo for valuable discussions.

REFERENCES

A.P.A (1994). DSM-IV Diagnostic and Statistical Manual of Mental Disorders, 4th Edition. Washington DC: American Psychiatric Association.

Auyeung B, Baron-Cohen S, Chapman E, Knickmeyer R, Taylor K, et al. (2006). Fetal testosterone and the Child Systemizing Quotient (SQ-C). European Journal of Endrocrinology 155,

Auyeung B, Baron-Cohen S, Chapman E, Knickmeyer R, Taylor K, et al. (2008). Fetal testosterone and autistic traits. British Journal of Psychology. online.

Auyeung, B., Baron-Cohen, S., Wheelwright, S., & Allison, C. (2008) The Autism Spectrum Quotient: Children's Version (AQ-Child) *Journal of Autism and Developmental Disorders*, XX, online.

Auyeung B, Baron-Cohen S, Chapman E, Knickmeyer R, Hackett G, 63 et al. (submitted) Fetal testosterone predicts sexually differentiated childhood behaviour in girls and boys.

Baron-Cohen S. (2002) The extreme male brain theory of autism. Trends in Cognitive Science 6, 248-254.

Baron-Cohen S. (2003) The Essential Difference: Men, Women and the Extreme Male Brain. London: Penguin.

Baron-Cohen S, & Wheelwright S. (2003) The Friendship 70 Questionnaire (FQ): An investigation of adults with Asperger 71 Syndrome or High Functioning Autism, and normal sex differences. Journal of Autism and Developmental Disorders 33, 73 509-517.

Baron-Cohen S, & Wheelwright S. (2004) The Empathy Quotient 75 (EQ). An investigation of adults with Asperger Syndrome or 76 High Functioning Autism, and normal sex differences. Journal of Autism and Developmental Disorders 34, 163-175.

Baron-Cohen S, Lutchmaya S, & Knickmeyer R. (2004) Prenatal testosterone in mind: Amniotic fluid studies. Cambridge, MA: 80 MIT/Bradford Books.

Baron-Cohen S, Knickmeyer R, & Belmonte MK. (2005) Sex differences in the brain: Implications for explaining autism. Science 310, 819-823.

Baron-Cohen S, Jolliffe T, Mortimore C, & Robertson M. (1997) 85 Another advanced test of theory of mind: Evidence from very 86 high functioning adults with autism or Asperger Syndrome. Journal of Child Psychology and Psychiatry 38, 813-822.

Baron-Cohen S, Hoekstra RA, Knickmeyer R, & Wheelwright S. (2006) The Autism-Spectrum Quotient (AQ)-Adolescent 90 version. Journal of Autism & Developmental Disorders 36, 343-350.

Baron-Cohen, S, Ring, H, Wheelwright, S, Bullmore, E, Brammer, M, 93 Simmons, A, & Williams, S. (1999) Social intelligence in the 94 normal and autistic brain: An fMRI study. European Journal of 95 Neuroscience, 11, 1891-1898.

Baron-Cohen S, Wheelwright S, Skinner R, Martin J, & Clubley E. 97 (2001) The Autism Spectrum Quotient (AQ): Evidence from Asperger Syndrome/High Functioning Autism, Males and 99 Females, Scientists and Mathematicians. Journal of Autism and Developmental Disorders 31, 5–17.

Baron-Cohen S, Wheelwright S, Scahill V, Lawson J, & Spong A. (2001) Are intuitive physics and intuitive psychology independent? Journal of Developmental and Learning Disorders 5, 47–78. 104

AU: List first six authors before et al. wherever "et al." is highlighted

> AU: Please provide website, date last accessed for all the online references.

60

65

66

68





62

67

68

71

72

76

85

100

109

110

112



994 **Autism Spectrum Disorders**

- Baron-Cohen S, Richler J, Bisarya D, Gurunathan N, & Wheelwright S. (2003) The Systemising Quotient (SQ): An investigation of adults with Asperger Syndrome or High Functioning Autism and 3 normal sex differences. Philosophical Transactions of the Royal Society 358, 361-374.
- Baron-Cohen S, Ring H, Chitnis X, Wheelwright S, Gregory L, 6 et al. (2006) fMRI of parents of children with Asperger Syndrome: A pilot study. Brain Cognition 61, 122-130.
- 9 Berenbaum S, & Hines M. (1992) Early androgens are related to childhood sex-typed toy preferences. Psychological Medicine 3, 10 203-206. 11
- Berenbaum SA. (1999) Effects of early androgens on sex-typed activi-12 ties and interests in adolescents with congenital adrenal hyper-13 plasia. Horm Behav 35(1), 102-110. 14
- Berenbaum SA, & Snyder E. (1995) Early hormonal influences 15 on childhood sex-typed activity and playmate preferences: 16 Implications for the development of sexual orientation. 17 Developmental Psychology 31, 31-42. 18
- Casella R, Maduro MR, Lipshultz LI, & Lamb DJ. (2001) Significance 19 of the polyglutamine tract polymorphism in the androgen 20 receptor. Urology 58, 651-656. 21
- Chakrabarti S, & Fombonne E. (2001) Pervasive Developmental 22 23 Disorders in pre-school children. Journal of the American Medical Association 285, 3093-3099. 24
- 25 Chapman E, Baron-Cohen S, Auyeung B, Knickmeyer R, Taylor K, et al. (2006) Fetal testosterone and empathy: Evidence from 26 the Empathy Quotient (EQ) and the "Reading the Mind in the 27 Eyes" Test.' Social Neuroscience 1, 135-148. 28
- Clark MM, & Galef BG. (1998) Effects of intraurine positioin on the 29 behaviour and genital morphology of litter=bearing rodents. 30 Developmental Neurology 14, 197–211. 31
- Collaer M, & Hines M. (1995) Human behavioural sex differences: 32 A role for gonadal hormones during early development? 33 Psychological Bulletin 118, 55-107. 34
- 35 Constantino JN, & Todd RD. (2005) Intergenerational transmission of subthreshold autistic traits in the general population. Biol 36 Psychiatry 57(6), 655-660. 37
- Cooke BM, Tabibnia G, & Breedlove SM. (1999) A brain sexual 38 dimorphism controlled by adult circulating androgens. 39 Proceedings for the National Academy of Sciences of the USA 40 96(13), 7538-7540. 41
- Courchesne E, Redcay E, & Kennedy DP. (2004) The autistic brain: 42 Birth through adulthood. Curr Opin Neurol 17(4), 489–496. 43
- Dawood MY. (1977) Hormones in amniotic fluid. Am J Obstet 44 Gynecol 128, 576-583. 45
- Ehrhardt AA, & Baker SW. (1974) Fetal androgens, human central 46 nervous system differentiation, and behavior sex differences. In: 47 Freidman RC, Richart RR, Van de Wiele RL, editors. Sex differ-48 ences in behavior. New York: Wiley. 49
- Fein D, Waterhouse L, Lucci D, Pennington B, & Humes M. (1985) 50 Handedness and cognitve functions in pervasive developmen-51 52 tal disorders. Journal of Autism and Developmental Disorders 15, 53 323-333.
- Finegan JA, Bartleman B, & Wong PY. (1989) A window for the study 54 of prenatal sex hormone influences on postnatal development. 55 J Genet Psychol 150(1), 101-112. 56
- 57 Fitch R, H, & Denenberg V. (1998) A role for ovarian hormones in sexual differentiation of the brain. Behavioural and Brain 58 59 Sciences 21, 311–352.
- Fombonne E. (2005) The changing epidemiology of autism. Journal 60 of Applied Research into Intellectual Disabilities 18, 281-294. 61

- Fuchs F, & Klopper A. (1983) Endocrinology of Pregnancy. Philadelphia: Harper & Row.
- Garn SM, Burdi AR, Babler WJ, & Stinson S. (1975) Early prenatal attainment of adult metacarpal-phalangeal rankings and proportions. Am J Phys Anthropol 43(3), 327-332.
- George M, Costa D, Kouris K, Ring H, & Ell P. (1992) Cerebral blood flow abnormalities in adults with infantile autism. Journal of Nervous and Mental Diseases 180, 413-417.
- Geschwind N, & Galaburda AM. (1985) Cerebral lateralization: 70 Biological mechanisms, associations and pathology. III. A hypothesis and a program for research. Archives of Neurology 42, 634-654.
- Giedd JN, Viatuzis AC, Hamburger SD, Lange N, Rajapakse JC, et al. (1996) Quantatitive MRI of the temporal lobe, amygdala and hippocampus in normal human development: Ages 4-18 years. The Journal of Comparative Neurology 366, 223-230.
- Gillberg C. (1983) Autistic children's hand preferences: Results from an epidemiological study of infantile autism. Psychiatry Res 10(1), 21-30.
- Golan O, Baron-Cohen S, & Hill J (2006) The Cambridge Mindreading (CAM) Face-Voice Battery: Testing complex emotion recognition in adults with and without Asperger syndrome. Journal of Autism and Developmental Disorders 36, 169-183.
- Goy RW. (1980) Sexual Differentiation of the Brain. Cambridge, MA: The MIT Press.
- Grimshaw G, Sitarenios G, & Finegan J. (1995) Mental rotation at 7 years: Relations with prenatal testosterone levels and spatial play experiences. Brain Cogn 29, 85-100.
- Grumbach MM, & Shaw EB. (1998) Further studies on the treatment of congenital adrenal hyperplasia with cortisone: IV. Effect of cortisone and compound B in infants with disturbed electrolyte 92 metabolism, by John F. Crigler Jr, MD, Samuel H. Silverman, MD, and Lawson Wilkins, MD, Pediatrics, 1952;10:397-413. Pediatrics 102, 215-221.
- Grumbach MM, Hughes IA, & Conte FA. (2003) Disorders of sex differentiation. In: Larsen PR, Kronenburg HM, Melmed S, Polansky KS, editors. Williams Textbook of Endocrinology. Philadelphia: Saunders.
- Hampson E, Rovet JF, & Altmann D. (1998) Spatial reasoning in children with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. Developmental Neuropsychology 14, 299-320.
- Hardan AY, Minshew NJ, Harenski K, & Keshavan MS. (2001) Posterior Fossa Magnetic Resonance Imaging in Autism. Journal of the American Academy of Child & Adolescent Psychiatry 40(6), 105 666-672. 106
- Herbert MR, Ziegler DA, Makris N, Filipek PA, Kemper TL, et al. (2004) Localization of white matter volume increase in autism and developmental language disorder. Ann Neurol 55(4), 530-540.
- Hines M. (2004) Brain Gender. Oxford & New York: Oxford 111 University Press.
- Hines M, Ahmed SF, & Hughes IA. (2003a) Psychological outcomes and gender-related development in complete androgen insensitivity syndrome. Archives of Sexual Behaviour 32, 93-101.
- Hines M, Brook C, & Conway GS. (2004) Androgen and psychosexual development: Core gender identity, sexual orientation 117 and recalled childhood gender role behavior in women and men with congenital adrenal hyperplasia (CAH). J Sex Res 41(1), 75-81.
- Hines M, Fane BA, Pasterski VL, Mathews GA, Conway GS, et al. (2003b) Spatial abilities following prenatal androgen 122





83

93

106

112

116



- abnormality: Targeting and mental rotations performance 1 in individuals with congenital adrenal hyperplasia. Psychoneuroendocrinology 28, 1010-1026.
- Hoekstra RA, Bartels M, Cath DC, & Boomsma DI. (2008) Factor 4 Structure, Reliability and Criterion Validity of the Autism-Spectrum Quotient (AQ): A Study in Dutch Population and Patient Groups. Journal of Autism and Developmental Disorders.
- Ingudomnukul E, Baron-Cohen S, Knickmeyer R, & Wheelwright S. (2007) Elevated rates of testosterone-related disorders in a 10 sample of women with autism spectrum conditions. Hormones and Behavior 51, 597-604. 11
- Jancke L, Staiger JF, Schlaug G, Huang Y, & Steinmetz H. (1997) 12 The relationship between corpus callosum size and forebrain 13 volume. Cereb Cortex 7(1), 48-56.
- Jolliffe T, & Baron-Cohen S. (1997) Are people with autism or 15 Asperger's Syndrome faster than normal on the Embedded 16 Figures Task? J Child Psychol Psychiatry 38, 527-534. 17
- Jost A. (1961) The role of fetal hormones in prenatal development. 18 Harvey Lectures 55, 201-226. 19
- Jost A. (1970) Hormonal factors in the sex differentiation of the mam-20 malian foetus. Philosophical Transactions of the Royal Society of 21 London 259(828), 119-130. 22
- 23 Jost A. (1972) A new look at the mechanisms controlling sex differentiation in mammals. Johns Hopkins Med Journal 130, 38-53. 24
- 25 Judd HL, Robinson JD, Young PE, & Jones OW. (1976) Amniotic fluid testosterone levels in midpregnancy. Obstet Gynecol 48(6), 26 27
- Kimura D. (1999) Sex and Cognition. Cambridge, MA: MIT Press. 28
- Kimura D, & Hampson E. (1994) Cognitive pattern in men and 29 women is influenced by fluctuations in sex hormones. Current 30 *Directions in Psychological Science 3*, 57–61. 31
- Knickmeyer R, Baron-Cohen S, Raggatt P, & Taylor K. (2005) Fetal 32 testosterone, social cognition, and restricted interests in children. 33 Journal of Child Psychology and Psychiatry 45, 1-13. 34
- 35 Knickmeyer R, Baron-Cohen S, Hoekstra R, & Wheelwright S. (2006) Age of menarche in females with autism spectrum conditions. 36 37 Developmental Medicine and Child Neurology 48, 1007–1008.
- Knickmeyer R, Baron-Cohen S, Raggatt P, Taylor K, & Hackett G. 38 (2006) Fetal testosterone and empathy. Horm Behav 49(3), 39 40
- Knickmeyer R, Baron-Cohen S, Fane BA, Wheelwright S, Mathews 41 GA, et al. (2006) Androgens and autistic traits: A study of indi-42 viduals with congenital adrenal hyperplasia. Hormones and 43 Behavior 50, 148-153. 44
- Lawson J, Baron-Cohen S, & Wheelwright S. (2004) Empathising 45 and systemising in adults with and without Asperger Syndrome. 46 J Autism Dev Disord 34, 301-310. 47
- 48 Lutchmaya S, Baron-Cohen S, & Raggatt P. (2002) Fetal testosterone and vocabulary size in 18- and 24-month-old infants. Infant 49 Behav Dev 24(4), 418-424. 50
- Lutchmaya S, Baron-Cohen S, & Raggatt P. (2002) Fetal testosterone 51 52 and eye contact in 12 month old infants. Infant Behav Dev 25, 53 327-335.
- 54 Mallin SR, & Walker FA. (1972) Effects of the XYY karyotype in one of two brothers with congenital adrenal hyperplasia. Clin Genet 55 56
- Manning J, Baron-Cohen S, Wheelwright S, & Sanders G. (2001) 57 Autism and the ratio between 2nd and 4th digit length. Dev Med 58 59 Child Neurol 43, 160–164.
- Manning JT, Martin S, Trivers RL, & Soler M. (2002) 2nd to 4th digit 60 ratio and offspring sex ratio. J Theor Biol 217(1), 93-95. 61

- McManus IC, Murray B, Doyle K, & Baron-Cohen S. (1992) Handedness in childhood autism shows a dissociation of skill and preference. Cortex 28, 373-381.
- Merke DP, Fields JD, Keil MF, Vaituzis AC, Chrousos GP, et al. (2005) Children with classic congenital adrenal hyperplasia 66 have decreased amygdala volume: Potential prenatal and postnatal hormone effects. The Journal of Clinical Endrochrinology & Metabolism 88, 1760-1765.
- Milne E, White S, Campbell R, Swettenham J, Hansen P, et al. (2006) Motion and Form Coherence Detection in Autistic Spectrum 71 Disorder: Relationship to Motor Control and 2:4 Digit Ratio. Journal of Autism and Developmental Disorders 36, 1–13.
- Money J, & Ehrhardt AA. (1972) Man and Woman, Boy and Girl. Baltimore: Johns Hopkins University Press.
- Money J, Schwartz M, & Lewis VG. (1984) Adult erotosexual status and fetal hormonal mASDulinization and demAS-Dulinization: 46,XX congenital virilizing adrenal hyperplasia and 46,XY androgen-insensitivity syndrome compared. Psychoneuroendocrinology 9, 405-414.
- Nagami M, McDonough P, Ellegood J, & Mahesh V. (1979) Maternal and amniotic fluid steroids throughout human pregnancy. American Journal of Obstetrics and Gynaecology 134,
- Nass R, Baker S, Speiser P, Virdis R, Balsamo A, et al. (1987) Hormones and handedness: Left-hand bias in female congenital adrenal hyperplasia patients. Neurology 37, 711-715.
- Nordenstrom A, Servin A, Bohlin G, Larsson A, & Wedell A. (2002) Sex-typed toy play behavior correlates with the degree of prenatal androgen exposure assessed by CYP21 genotype in girls with congenital adrenal hyperplasia. Journal of Clinical Endocrinology & Metabolism 87, 5119-5124.
- Peters M. (1991) Sex, handedness, mathematical ability, and biological causation. Can I Psychol 45(3), 415-419.
- Phoenix CH, Goy RW, Gerall AA, & Young WC. (1959) Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. Endocrinology 65, 369-382.
- Quadagno DM, Briscoe R, & Quadagno JS. (1977) Effect of perinatal gonadal hormones on selected nonsexual behavior patterns: 100 A critical assessment of the nonhuman and human literature. 101 Psychological Bulletin 84, 62-80.
- Resnick S, Berenbaum S, Gottesman I, & Bouchard T. (1986) Early hormonal influences on cognitive functioning in congenital adrenal hyperplasia. Developmental Psychology 22, 191-198.
- Ring H, Baron-Cohen S, Williams S, Wheelwright S, Bullmore E, Brammer M, & Andrew C. (1999) Cerebral correlates of preserved cognitive skills in autism. A functional MRI study of Embedded Figures Task performance. *Brain*, 122, 1305–1315.
- Robinson JD, Judd HL, Young PE, Jones OW, & Yen SS. (1977) 110 Amniotic fluid androgens and estrogens in midgestation. J Clin 111 Endocrinol Metab 45(4), 755-761.
- Rohde Parfet KA, Ganjam VK, Lamberson WR, Rieke AR, Vom Saal F, 113 et al. (1990) Intrauterine position effects in female swine: 114 Subsequent reproductive performance and social and sexual behaviour. Applied Animal Behaviour Science 26: 349-362.
- Rommerts FFGHK, Gromoll J, Cato ACB, Hiort O, Zitzmann M, Christiansen K, et al. (2001) Testosterone: Action, deficiency, substitution; Nieschlag E, Behre H, editors. Cambridge: Cambridge University Press.
- Satz P, Soper H, Orsini D, Henry R, & Zvi J. (1985) Handedness 121 subtypes in autism. Psychiatric Annals 15, 447-451. 122







996 **Autism Spectrum Disorders**

- Schindler AE. (1982) Hormones in human amniotic fluid. Monogr Endocrinol 21, 1-158.
- Scott F, Baron-Cohen S, Bolton P, & Brayne C. (2002) Prevalence of autism spectrum conditions in children aged 5-11 years in Cambridgeshire, UK. Autism 6(3), 231–237.
- Scott F, Baron-Cohen S, Bolton P, & Brayne C. (2002) The CAST (Childhood Asperger Syndrome Test): Preliminary development of UK screen for mainstream primary-school children. Autism 6(1), 9–31.
- 10 Shah A, & Frith U. (1983) An islet of ability in autism: A research note. Journal of Child Psychology and Psychiatry 24, 613-620. 11
- Smail PJ, Reyes FI, Winter JSD, & Fairman C. (1981) The fetal hor-12 monal environment and its effect on the morphogenesis of the genital system. In: Kogan SJ, Hafez ESE, editors. Pediatric andrology (pp. 9–19). The Hague: Martinus Nijhoff.
 - Soper H, Satz P, Orsini D, henry R, Zvi J, et al. (1986) Handedness patterns in autism suggests subytpes. Journal of Autism and Developmental Disorders 16, 155-167.
 - Sparks BF, Friedman SD, Shaw DW, Aylward EH, Echelard D, et al. (2002) Brain structural abnormalities in young children with autism spectrum disorder. Neurology 59, 184-192.
- Stodgell CJ, Ingram JI, & Hyman SL. (2001) The role of candidate 22 23 genes in unravelling the genetics of autism. International Review of Research in Mental Retardation 23, 57-81. 24
- 25 Tordjman A, Ferrari P, Sulmont V, Duyme M, & Roubertoux P. (1997) Androgenic Activity in Autism. American Journal of 26 Psychriatry 154, 11. 27

- Tulchinsky D, & Little AB. (1994) Maternal-fetal Endocrinology. Philadelphia & London: W B Saunders.
- Wakabayashi A, Baron-Cohen S, & Wheelwright S. (2004) The Autism 30 Spectrum Quotient (AQ) Japanese version: Evidence from highfunctioning clinical group and normal adults. Japanese Journal 32 of Psychology 75, 78-84.
- Wakabayashi A, Baron-Cohen S, Wheelwright S, & Tojo Y. (2006) 34 The Autism-Spectrum Quotient (AQ) in Japan: A cross-cultural 35 comparison. Journal of Autism & Developmental Disorders 36, 36 263-270.
- Wakabayashi A, Baron-Cohen S, Uchiyama T, Yoshida Y, Tojo Y, et al. (2007) The Autism-Spectrum Quotient (AQ) Children's Version in Japan: A Cross-Cultural Comparison. Journal of Autism and Developmental Disorders 37, 491–500.
- Wheelwright S, Baron-Cohen S, Goldenfeld N, Delaney J, Fine D, 42 et al. (2006) Predicting Autism Spectrum Quotient (AQ) 43 from the Systemizing Quotient-Revised (SQ-R) and Empathy 44 Quotient (EQ). Brain Research 1079, 47-56.
- Williams J, Scott F, Allison C, Bolton P, Baron- Cohen S, et al. (2005) The CAST (Childhood Asperger Syndrome Test): Test accuracy. Autism, 45-68.
- Williams JG, Allison C, Scott FJ, Bolton PF, Baron-Cohen S, et al. (2008) The Childhood Autism Spectrum Test (CAST): Sex Differences. Journal of Autism and Developmental Disorders.



13

14

15

16

17

18

19

20

21

Verify

lower

case 'h'



The following need to be added to the reference list:

Mathews GA, Fane BA, Conway GS, Brook CG, Hines M (2009) Personality and congenital adrenal hyperplasia: possible effects of prenatal androgen exposure. Hormones and Behavior 55: 285-291. Helleday J, Edman G, Ritzen EM, Siwers B (1993) Personality characteristics and platelet MAO activity in women with congenital adrenal hyperplasia (CAH). Psychoneuroendocrinology 18: 343-354.

Resnick SM (1982) Psychological functioning in individuals with congenital adrenal hyperplasia: early hormonal influences on cognition and personality. Minneapolis: University of Minnesota. Kuhnle U, Bullinger M (1997) Outcome of congenital adrenal hyperplasia. Pediatric Surgery International 12: 511-515.